



Pest Alert


Emerald Ash Borer

(*Agrilus planipennis*)



Larva

D-Shaped Emergence Hole




Adult



Actual Size
8.5 – 14mm

G.G.Wood, CFIA



Tree Decline

James W. Smith, MDA



Ash Leaf

Ken Marchant, CFIA



S-Shaped Tunnels

CFIA Windsor

The Emerald Ash Borer (EAB) is an insect pest that is highly destructive to ash trees. Native to eastern Asia, this pest was first discovered in Canada and the U.S. in 2002. The EAB has killed millions of ash trees in Manitoba, Ontario, Quebec and the United States, and poses a major economic and environmental threat to urban and forested areas in both countries. The EAB attacks and kills all species of ash (except Mountain ash which is not a true ash).

How does the EAB spread?

While the EAB can fly up to several kilometres, another significant factor contributing to its spread is the movement of firewood, nursery stock, trees, logs, lumber, wood with bark attached and wood or bark chips.

Signs of infestation

Tree decline, including:

- thinning crown
- diminished density of leaves
- evidence of adult beetle feeding on leaves
- long shoots growing from the trunk or branches
- vertical cracks in the trunk
- small D-shaped emergence holes
- S-shaped tunnels under the bark filled with fine sawdust

Contact the CFIA

If you see suspected signs of infestation on your ash trees or if you plan on moving firewood (e.g., when camping, relocating or managing a woodlot), contact the Canadian Food Inspection Agency (CFIA) for more information.

inspection.gc.ca/pests

Help Protect Canada's Trees and Forests.

