



# STOP THE SPREAD

## Emerald ash borer (*Agrilus planipennis*)

**The emerald ash borer (EAB)** is a highly destructive insect that targets ash trees. Native to eastern Asia, it was first discovered in Canada and the United States (US) in 2002. Since then, it has killed millions of ash trees in eastern Canada and the US, posing a significant economic and environmental threat to both urban and forested areas. This pest kills all species of ash (except for mountain ash *Sorbus* spp., which is not a true ash). EAB has been detected in 36 US states and six Canadian provinces.

### HOW DOES IT SPREAD?



While EAB can fly up to 10 kilometres, another significant factor contributing to its spread is the movement of firewood, nursery stock, trees, logs, lumber, wood with bark attached and wood or bark chips.

### WHAT TO LOOK FOR - SIGNS AND SYMPTOMS OF INFESTATION



Thinning crown (diminished density of branches and leaves)



Epicormic shoots (i.e. water sprouts) growing from the trunk or branches




Small D-shaped emergence holes



S-shaped tunnels under the bark filled with fine sawdust



Feeding damage on leaves caused by the beetle



Vertical cracks in the trunk



Contact the Canadian Food Inspection Agency if you see suspicious signs or symptoms of emerald ash borer on your ash.

Help protect Canada's trees and forests. It's up to all of us to prevent the spread of invasive species.



Photo Credits: T. Kimoto (CFIA), Dr. James E. Zaboltny (USDA, APHIS), Debbie Miller (USDA Forest Service)

© His Majesty the King in Right of Canada, as represented by the Minister of Health, 2025.  
Catalogue No.: A104-643/2025E-PDF ISBN: 978-0-660-76436-8  
Aussi disponible en français.



Canadian Food  
Inspection Agency

Agence canadienne  
d'inspection des aliments

Canada